

A Multi-Wavelength Mini Lidar for Measurements of Marine Boundary Layer Aerosol and Water Vapor Fields

Shiv K. Sharma

Hawaii Institute of Geophysics & Planetology
2525 Correa Rd., Honolulu HI 96822
phone: (808) 956-8476 fax: (808) 956-3188
email: sksharma@soest.hawaii.edu

Barry R. Lienert

Hawaii Institute of Geophysics & Planetology
phone: (808) 956-7815 fax: (808) 956-3188
email: lienert@soest.hawaii.edu

John N. Porter

Hawaii Institute of Geophysics & Planetology
phone: (808) 956-6483 fax: (808) 956-3188
email: jporter@soest.hawaii.edu

Award #: N000149610317

<http://www.soest.hawaii.edu/lidar>

LONG-TERM GOALS

Our long-term goal is to improve our understanding of dynamics of marine aerosols and water vapor fields in the coastal marine boundary layer using a scanning lidar and meteorological parameters.

OBJECTIVES

Our scientific objectives are to collect well-calibrated lidar data sets that can be used to improve and develop models of the aerosol optical properties in the coastal marine boundary layer (MBL). Various aerosol models exist (e.g., Fitzgerald, 1989), but few are appropriate for coastal regions. We are studying the vertical aerosol structure in the 15-m of the atmosphere directly above the ocean surface.

APPROACH

We are using a scanning multi-wavelength lidar to measure the 4-D (space and time) aerosol optical fields in order to characterize the aerosol properties in a marine setting (Sharma et al., 2001). These measurements have been carried out at Bellows Air Force (AFS) next to the University of Hawaii's Meteorological Tower (21° 21.848' N, 157° 42.584' W). We are able to study the spatial structures of the aerosol scattering fields out to distances of up to 10 km from the shore. The scanning lidar data enables us to place the aerosol characteristics being observed by the shore-based instruments in the context of much larger scale variability in the aerosol scattering fields. Dr. Shiv Sharma is the project director involved in all aspects of these efforts. Dr. Barry Lienert has developed the software and supervises the data collection. Dr. John Porter is involved in calibration and modeling of the data.

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2002		2. REPORT TYPE		3. DATES COVERED 00-00-2002 to 00-00-2002	
4. TITLE AND SUBTITLE A Multi-Wavelength Mini Lidar for Measurements of Marine Boundary Layer Aerosol and Water Vapor Fields				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Hawaii Institute of Geophysics & Planetology,,2525 Correa Rd.,,Honolulu,,HI, 96822				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT Our long-term goal is to improve our understanding of dynamics of marine aerosols and water vapor fields in the coastal marine boundary layer using a scanning lidar and meteorological parameters.					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 9	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

WORK COMPLETED DURING 2002

- (1) We have processed lidar and meteorology data during the SEAS experiment to develop a better understanding of the coastal optical properties at Bellows beach (Porter et al., 2002).
- (2) We have completed a preliminary study of the relationship between plume height over a reef with wind speed (Sharma et al., 2002)
- (3) We have derived reliable sea salt size distributions from genetic inversion of polar nephelometer data (Lienert et al, 2002).

RESULTS




Figure 1. Height of the top of the mixed layer derived from lidar 2-D scans (black dots) and Lihue, Kauai National Weather Service sonde data (blue circles). The height of the trade wind inversion base and top (from the Lihue soundings) are also shown in the upper portion of the figure. The mixed layer height varies between 300 and 1300 m, while the trade wind inversion height varies between 1400 and 2300 m, with an average thickness of about 300 m. No obvious correlation is apparent between the mixed layer and trade wind inversion heights.

During the SEAS experiment (4/19/2000-4/23/2000) different investigators brought their instruments together for a focused study of coastal marine aerosols. During this period the spatial and temporal distribution of coastal aerosols was measured by the University of Hawaii scanning lidar (Sharma et al., 2001) providing several interesting features of coastal marine aerosols. One of the prominent

features which is evident in the lidar data is a mixed layer from the surface up to approximately 500 m height. It is important to understand the physics of this layer in order to explain processes occurring in the lowest 15 m (Kepert and Fairall, 1999). Models of the mixed layer thickness have often portrayed it as gradually varying as the air mass is advected. Our lidar data during the SEAS experiments suggested that it actually varied significantly over short temporal and spatial scales. This is illustrated in Fig. 1 which shows the height of the mixed layer derived from the lidar data as well as from soundings (on Lihue, Kauai) during the SEAS experiment. The height of the base and top of the trade wind inversion is also shown for the Kauai soundings. The rapid variability suggests cloud scale processes may be responsible for these variations. Further studies are needed to investigate these rapid variations.




Figure 2. Spatial distribution of lognormal average aerosol scattering coefficient derived from the scanning lidar (for the whole SEAS lidar data set). The beach is a range zero (on the right) and the outer reef is at 2 km range. The left side of the image is upwind of the lidar site. The extinction in the 500 m thick mixed layer ($\sim 2 \times 10^{-5} \text{ m}^{-1}$) increases by a factor of about three at distances less than 2 km from the coastline. The 30 m thick region at the surface, 1.5–2.2 km from the coast, is due to spray from waves breaking on reefs.




Figure 3. Same as Figure 3 but showing the lognormal standard deviation. Standard deviations in the mixed layer are $1-2 \times 10^{-5} \text{ m}^{-1}$, increasing to $4 \times 10^{-5} \text{ m}^{-1}$ at heights of 400-800 m and to 1×10^{-4} over the reef. The larger variability in the bottom left of the image is an artifact of poor signal to noise. This has since been improved through the use of a logarithmic amplifier.

We studied the statistical variability of the aerosol fields at Bellows over a five day period during the SEAS experiment (Porter et al., 2002). We have calculated the lognormal average and the lognormal standard deviation of aerosol scattering coefficients in rectangular boxes spanning a complete vertical section (Figs. 2 and 3). It is evident that island blocking is causing enhanced scatter (orange region in Fig. 2) to develop 2 km from the coastline. This region is not apparent in the standard deviation plot in Fig. 3, indicating that is a persistent feature in the offshore aerosol. Visual observations during SEAS suggest that is related to cloud development. The enhanced scatter persists all the way down to sea level, possibly as a result of cloud-related drizzle or virga. The spray from breaking waves over the reefs (red region near the surface at 1.3-2.2 km range) remains below 30 m for the trade wind conditions during SEAS ($\sim 7 \text{ m/s}$). Large standard deviations in this region (Fig. 3) indicate that the reef spray scatter varied episodically with tide level and wave height.




Figure 4. Vertical distribution of aerosol extinction over the waves breaking on Bellows Beach (squares) and the those obtained during the EOPACE experiment over the waves at Scripps Pier, San Diego , California (circles). The extinction at Bellows increases from $5 \times 10^{-5} \text{ m}^{-1}$ at heights greater than 7 m to $6 \times 10^{-4} \text{ m}^{-1}$ at a height of 1m. The EOPACE extinctions increase from $7 \times 10^{-5} \text{ m}^{-1}$ at a height of 10 m to $4 \times 10^{-4} \text{ m}^{-1}$ at 4 m, the lowest height measured.

A vertical profile of extinction (532 nm) directly over waves breaking on the beach was made from lidar data collected on 6/11/99 and is shown in Fig. 4. The extinctions measured during the EOPACE (Electro-optical Propagation Assessment in Coastal Environments) experiment are also shown. During the EOPACE experiment the spray concentrations were larger possibly due to the lighter winds ($\sim 2 \text{ m/s}$) versus the 4.7 m/s winds during the Bellows Beach measurements. The waves at Scripps Pier are also generally larger than those at Bellows Beach. Assuming a wind speed of 5 m/s results in an average salt mass flux of $807 \mu\text{g m}^{-2} \text{ sec}^{-1}$ from Bellows Beach waves which is within the range of $4562\text{--}1034 \mu\text{g m}^{-2} \text{ sec}^{-1}$ reported by de Leeuw et al., (2000) for breaking waves at Scripps Pier, California.

During the SEAS experiment we also measured the aerosol phase function (at 532 nm) with a polar nephelometer (Porter et al., 1998). Inversion of the polar nephelometer data (Lienert et al., 2002) resulted in a coarse mode number size distribution (center wet diameter $\sim 1.8 \mu\text{m}$) which was in agreement with inversions of our lidar data (Shifrin et al., 2002) as well as direct measurements (Clarke et al., 2002) of the aerosol size distribution after inlet loss corrections. Fig. 5 shows the measured phase function (due to aerosol and molecular scatter) along with the total phase function calculated from the aerosol inversions. By successfully fitting this data in the angular range $140\text{--}180^\circ$, we are able to place much better constraints on the coarse mode size distribution than could be achieved with lidar measurements at 532 and 1064 nm wavelength.




Figure 5. *Measured and predicted (for bimodal and unimodal inverted lognormal aerosol models) total (aerosol plus molecular) angular scattering 5m above sea level on Bellows Beach. The poor fit for scattering angles less than 30° (data is 20% lower than theoretical) is due to improper calibration of an attenuation filter which was used in the early version of the polar nephelometer. The majority of the remainder of the data is within 15% of the theoretical curves.*

TRANSITIONS

We are presently carrying out further studies of plume height and spatial distribution as a function of wind speed. These studies will be carried out using the extensive lidar data set collected at Bellows from 1999-2002. We have already shown that wavelengths below $\sim 1\ \mu\text{m}$ are not sufficient to resolve the coarse mode sea salt (Lienert et al., 2001). However, we have found that the addition of a $9.25\ \mu\text{m}$ lidar channel dramatically improves the constraints on the coarse mode size distribution. Further work in this area is planned.

RELATED PROJECTS

Our lidar efforts are closely related to the work being carried out by Dr. Antony Clarke. We are comparing our lidar measurements with his *in situ* aerosol size distribution measurements as well as our polar nephelometer inversion results. These independent constraints on the aerosol size distributions at Bellows form an essential part of our ongoing efforts to interpret the high-quality multi-wavelength lidar data we acquired in 2001. We are also continuing our cooperation with Dr. Kuseil Shifrin's group.

REFERENCES

- De Leeuw, G., F.P. Neele, Hill, M., M. H. Smith, and E. Vignati, 2000: Production of sea spray aerosol in the surf zone, *J. Geophys. Res.*, **105**, 29397-29409.
- Fitzgerald, J.W., 1989, Model of the aerosol extinction profile in a well-mixed marine boundary layer, *Applied Optics*, 28, 3534-3538.
- Keprt, J.D. and C.W. Fairall, 1999. Modelling the interaction between the atmospheric boundary layer and evaporating sea spray droplets, *Third Conference on Boundary Layers and Turbulence, Dallas TX, Jan 10-15, 1999* (http://www.bom.gov.au/bmrc/daas/da_staff/keprt/sea_spray.htm)
- B. R. Lienert, J.N. Porter and S.K. Sharma, 2001. Repetitive Genetic Inversion of Optical Extinction Data, *Appl. Optics*, **40**, 3476-3482.
- Lienert, B.R., J. N. Porter and S. K. Sharma, 2002. Aerosol Size Distributions from Genetic Inversion of Polar Nephelometer Data (*submitted to J. Atmospheric and Oceanic Tech.*)
- Porter, J.N., S.K. Sharma, B. R. Lienert, and E. Lau, 2002. Aerosol Scattering Fields Over Bellows Beach, Oahu During the SEAS Experiment, (*submitted to J. Atmospheric and Oceanic Tech.*).
- Porter, J.N., T.F. Cooney, and C. Motell, 1998: Coastal Aerosol Phase Function Measurements With A Custom Polar Nephelometer, *Ocean Optics XIV Conference, Kona, Hawaii*, pp. 5.
- Sharma, S. K., B. R. Lienert and J. N. Porter, 2001. Scanning Lidar Measurements of Marine Aerosol Fields At A Coastal Lidar Site in Hawaii. *Proceedings of the SPIE Lidar Remote Sensing for Industry and Environment Monitoring*, Sendai, Japan, SPIE vol. **1453**, 159-166.
- Sharma, S. K., B. R. Lienert and J. N. Porter, 2002. Multi-wavelength Scanning Lidar Measurements of the Effect of Wind Speed on Marine Aerosol Fields Generated by Breaking Waves, *Proceedings of the SPIE Conference on Ocean Remote Sensing and Applications, Hangzhou, China, 23-27 October, 2002*.
- Shifrin, K.S. and I. G. Zolotov, 2002. The Use of Direct Observations Over the Aerosol Particle Size Distribution for Inverting Lidar Data (*submitted to J. Atmospheric and Oceanic Tech.*).

PUBLICATIONS

- B. Lienert, J. Porter, N. Ahlquist, D. Harris, S. Sharma, A 50-MHz logarithmic amplifier for use in lidar measurements, *J. Atmos. Ocean. Tech.*, 19, 654-657, 2002.
- Porter, J. N.; Horton, K. A.; Mougini-Mark, P. J.; Lienert, B.; Sharma, S. K.; Lau, E.; Sutton, A. J.; Elias, T.; Oppenheimer, C., 2002. Sun photometer and lidar measurements of the plume from the Hawaii Kilauea Volcano Pu'u O'o vent: Aerosol flux and SO₂ lifetime, *Geophys. Res. Lett.*, *in press*
- A. Clarke, V. Kapustin, S. Howell, K. Moore, B. Lienert, S. Masonis, T. Anderson, D. Covert, K. Shifrin, I. Zolotov, 2002. Sea-salt size-distributions from breaking waves: implications for marine aerosol production and optical extinction measurements during SEAS (*submitted to J. Atmos. Ocean. Tech.*)
- Lienert, B.R., J. N. Porter and S. K. Sharma, 2002. Aerosol Size Distributions from Genetic Inversion of Polar Nephelometer Data (*submitted to J. Atmos. Ocean. Tech.*)
- Porter, J.N., S.K. Sharma, B. R. Lienert, and E. Lau, Aerosol Scattering Fields Over Bellows Beach, Oahu During the SEAS Experiment, 2002. (*submitted to J. Atmos. and Ocean. Tech.*)

Porter, J.N., B. Lienert, S. K. Sharma, and H. W. Hubble, 2002. A Low Cost Portable Lidar System To Measure Aerosol Optical Properties., *J. Atmos. and Ocean. Tech.*, *in press*.